VOL. 506 JOURNAL OF CHROMATOGRAPHY MAY 11, 1990

CONTENTS

13TH INTERNATIONAL SYMPOSIUM ON COLUMN LIQUID CHROMATOGRAPHY, STOCKHOLM, JUNE 25–30, 1990, PART I

Foreword
by D. Westerlund 1

THEORY
Retention mechanisms

Correlation between chemical structure of non-congeneric solutes and their retention on polybutadiene-coated alumina
by R. Kaliszan and K. Ośmiałowski (Gdańsk, Poland) 3

High-performance liquid chromatography of amino acids, peptides and proteins. XCVII. The influence of the gradient elution mode and displacer salt type on the retention properties of closely related protein variants separated by high-performance anion-exchange chromatography
by A. N. Hodder, M. I. Aguilar and M. T. W. Hearn (Clayton, Australia) 17

Application of chromatographic retention data in an investigation of a quantitative structure–nucleotide incorporation rate relationship
by K. Valkó, T. Cserháti, I. Fellegvári, J. Sági and S. Szemzö (Budapest, Hungary) . . . 35

Secondary equilibria and their interaction with chromatographic transport
by H. Poppe (Amsterdam, The Netherlands) 45

Studies on system peaks in ion-pair adsorption chromatography. III. Regulation of system peak gradient retention for obtaining analyte peak compression
by T. Fornstedt, D. Westerlund and A. Sokolowski (Uppsala, Sweden) 61

Multiple peak formation in reversed-phase liquid chromatography of ramipril and ramiprilate
by S. Gustafsson, B.-M. Eriksson and I. Nilsson (Mölndal, Sweden) 75

Extension of the electrostatic retention model of reversed-phase ion-pair chromatography to include the simultaneous effects of the organic modifier and the pairing ion
by A. Bartha and Gy. Vigh (College Station, TX, U.S.A.) and J. Ståhlberg (Sjödertälje, Sweden) 85

Silanol effects in reversed-phase liquid chromatography
by T. Welsch and H. Frank (Leipzig, G.D.R.) and Gy. Vigh (College Station, TX, U.S.A.) 97

Reversed-phase high-performance liquid chromatography in the investigation of the hydrophobicity of selected ketones
by A. Siwek and J. Śliwiok (Katowice, Poland) 109

Partition coefficients and capacity factors of some nucleoside analogues
by A. P. Cheung and D. Kenney (Menlo Park, CA, U.S.A.) 119

Application of a computer-assisted high-performance liquid chromatographic multi-wavelength ultraviolet detection system to simultaneous toxicological drug analyses
by M. Hayashida, M. Nihira and T. Watanabe (Tokyo, Japan) and K. Jinno (Toyohashi, Japan) 133

Chromatographic characteristics of nucleic acid components on vinyl alcohol copolymer gel columns
by K. Noguchi, Y. Yanagihara, M. Kasai and N. Hirata (Kawasaki, Japan) 145

Retention behaviour of mono- and dicarboxylic acids, carbohydrates and alcohols in ion-exclusion chromatography
by E. Papp (Veszprém, Hungary) and P. Keresztes (Budapest, Hungary) 157

Chemometrics and chromatography

Optimization of chromatographic methods by a combination of optimization software and expert systems
by P. J. Schoenmakers (Eindhoven, The Netherlands), A. Peeters (Brussels, Belgium) and R. J. Lynch (Cambridge, U. K.) 169

Theoretical optimization of open-tubular columns for liquid chromatography with respect to mass loadability
by P. P. H. Tock, P. P. E. Duijsters, J. C. Kraak and H. Poppe (Amsterdam, The Netherlands) 185

Selective determination of trace levels of phenol in river water using electrochemical concentration modulation correlation chromatography
by M. Engelsma, W. Th. Kok and H. C. Smit (Amsterdam, The Netherlands) 201

Validation of an expert system for the selection of initial high-performance liquid chromatographic conditions for the analysis of basic drugs
by F. Maris, R. Hindriks and J. Vink (Oss, The Netherlands) and A. Peeters, N. Vanden Driessche and L. Massart (Brussels, Belgium) 211

Resolution of unresolved peaks containing unknown components by high-performance liquid chromatography with multi-wavelength detection
by I. Sakuma (Saitama, Japan), N. Takai and T. Dohi (Tokyo, Japan), Y. Fukui (Saitama, Japan) and A. Ohkubo (Tokyo, Japan) 223

Optimization of mobile phase composition for high-performance liquid chromatographic separation by means of the overlapping resolution mapping scheme
by S. F. Y. Li, H. K. Lee and C. P. Ong (Singapore, Singapore) 245

Regulation of peak compression effects for substituted benzamides in reversed-phase liquid chromatography
by L. B. Nilsson (Södertälje, Sweden) 253

Towards a chromatographic quantitator
by P. A. Bristow (Macclesfield, U.K.) 265

High-performance liquid chromatographic determination of guanidino compounds by automated pre-column fluorescence derivatization
by V. K. Boppana and G. R. Rhodes (King of Prussia, PA, U.S.A.) 279

Multiple absorbance ratio correlation—a new approach for assessing peak purity in liquid chromatography
by J. G. D. Marr, G. G. R. Seaton, B. J. Clark and A. F. Fell (Bradford, U.K.) 289

TECHNIQUES

Stationary phases

Aligned fiber columns for size-exclusion chromatography
by M. Czok and G. Guiochon (Knoxville and Oak Ridge, TN, U.S.A.) 303

High-performance liquid chromatography of transfer ribonucleic acids on spherical hydroxyapatite beads
by Y. Yamakawa, K. Miyasaka and T. Ishikawa (Tokyo, Japan), Y. Yamada (Tochigi, Japan) and T. Okuyama (Tokyo, Japan) 319

Peptide behaviour and analysis on a chemically stable C_{18}-bonded vinyl alcohol copolymer column with alkaline and acidic eluents
by T. Uchida, T. Ohtani, M. Kasai, Y. Yanagihara and K. Noguchi (Kawasaki, Japan) and H. Izu and S. Hara (Osaka, Japan) 327

Retention times and heats of adsorption of aromatic compounds on carbon adsorbents
by S. A. Eltekova (Moscow, U.S.S.R.) 335

Investigations by ^{29}Si cross-polarization magic angle spinning NMR spectroscopy of reaction pathways of silica gel polyfunctional modification
by B. Pfleiderer, K. Albert and E. Bayer (Tübingen, F.R.G.) 343

Novel, highly deactivated reversed-phase for basic compounds
by T. L. Ascah and B. Feibush (Bellefonte, PA, U.S.A.) 357

Development and use of carbon adsorbents in the liquid chromatographic separation of isomers
by F. Belliardo and O. Chiantore (Turin, Italy), D. Berek and I. Novàk (Bratislava, Czechoslovakia) and C. Lucarelli (Rome, Italy) 371

Lipid-vesicle-surface chromatography
by Q. Yang, M. Wallstén and P. Lundahl (Uppsala, Sweden) 379

Effect of polystyrene coating on pore, structural and chromatographic properties of silica packings
by A. Kurganov, O. Kuzmenko and V. A. Davankov (Moscow, U.S.S.R.) and B. Eray, K. K. Unger and U. Trüdinger (Mainz, F.R.G.) 391

Methods

Indirect fluorometric detection in open-tubular capillary column chromatography
by W. D. Pfeffer and E. S. Yeung (Ames, IA, U.S.A.) 401

Application of low-angle laser light scattering detection in the field of biochemistry. Review of recent progress
by T. Takagi (Osaka, Japan) 409

Determination of amoxicillin in plasma by high-performance liquid chromatography with fluorescence detection after on-line oxidation
by H. Mascher and C. Kikuta (Vienna, Austria) 417

Selective post-column liquid chromatographic determination of sugars in spent sulphite liquor with two enzymatic electrochemical detectors in parallel
by G. Marko-Varga (Lund, Sweden), E. Dominguez (Madrid, Spain) and B. Hahn-Hägerdal and L. Gorton (Lund, Sweden) 423

Coupling of microcolumn high-performance liquid chromatography with Fourier transform infrared spectrometry
by K. Jinno and C. Fujimoto (Toyohashi, Japan) 443

In vivo microdialysis sampling coupled to liquid chromatography for the study of acetaminophen metabolism
by D. O. Scott, L. R. Sorensen and C. E. Lunte (Lawrence, KS, U.S.A.) 461

Determination of water-soluble vitamins in blood and plasma by coupled-column liquid chromatography
by K. Johansen and P. O. Edlund (Solna, Sweden) 471

Displacement chromatography on cyclodextrin–silicas. III. Enantiomer separations
by Gy. Vigh, G. Quintero and Gy. Farkas (College Station, TX, U.S.A.) 481

Solutions of the equilibrium and semi-equilibrium models of chromatography
by S. Golshan-Shirazi and G. Guiochon (Knoxville, TN, U.S.A.) 495

Hydrodynamic chromatography of macromolecules on small spherical non-porous silica particles
by G. Stegeman, R. Oostervink, J. C. Kraak and H. Poppe (Amsterdam, The Netherlands) and K. K. Unger (Mainz, F.R.G.) 547

Comparison of stationary phases for packed-column supercritical fluid chromatography
by P. J. Schoenmakers, L. G. M. Uunk and H.-G. Janssen (Eindhoven, The Netherlands) 563

Chiral separations

Chiral separations of atropine and homatropine on an α_1-acid glycoprotein-bonded stationary phase
by E. Arvidsson and S. O. Jansson (Mölndal, Sweden) and G. Schill (Uppsala, Sweden) . 579

Chiral chloroformates as transparent reagents for the resolution of metoprolol enantiomers by reversed-phase liquid chromatography
by M. Ahnoff, S. Chen, A. Green and I. Grundevik (Mölndal, Sweden) 593

Direct high-performance liquid chromatographic determination of the enantiomers of alfuzosin in plasma on a second-generation of α_1-acid glycoprotein chiral stationary phase
by A. Rouchouse, M. Manoha, A. Durand and J. P. Thenot (Paris, France) 601

Separation of the enantiomeric intermediates of some platelet-activating factor analogues on a naphthylalanine-type Pirkle column
by P. L. Camacho, E. Geiger and Gy. Vigh (College Station, TX, U.S.A.) and R. Webster and D. H. Thompson (Beaverton, OR, U.S.A.) 611

High-performance liquid chromatographic resolution of racemic 1,4-benzodiazepin-2-ones by means of a β-cyclodextrin silica bonded chiral stationary phase
by C. Bertucci, E. Domenici, G. Uccello-Barretta and P. Salvadori (Pisa, Italy) 617

Direct high-performance liquid chromatographic separation of (+)- and (−)-medetomidine hydrochloride with an α_1-acid glycoprotein chiral column
by G. Örn, K. Lahtonen and H. Jalonen (Turku, Finland) 627

Author Index 637

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-printed asterisk (*)